

Singularity confinement and algebraic integrability

S. Lafortune^{a)} and A. Goriely^{b)}

Department of Mathematics, University of Arizona, Tucson, Arizona 85721-0089

(Received 11 October 2003; accepted 12 November 2003)

Two important notions of integrability for discrete mappings, algebraic integrability and singularity confinement, have been used for discrete mappings. Algebraic integrability is related to the existence of sufficiently many conserved quantities and singularity confinement is associated with the local analysis of singularities. In this article, the relationship between these two notions is explored for birational autonomous mappings. The main result of this article is that algebraically integrable mappings are shown to have the singularity confinement property. Using this result, the proof of the nonexistence of algebraic conserved quantities for a class of discrete systems is given. © 2004 American Institute of Physics.

[DOI: 10.1063/1.1640797]

I. INTRODUCTION

One of the first notions of integrability was introduced by Liouville in the 19th century for Hamiltonian systems in classical mechanics. This type of integrability is based on quantities whose values do not change in time, the so-called *constants of the motion* or *first integrals*. For instance, in many mechanical systems the total energy and the linear and angular momenta are conserved through the dynamics and are examples of such constants. Liouville's fundamental contribution was to prove that if a given Hamiltonian system admits enough first integrals, it can be solved explicitly by quadratures. More precisely, a $2n$ -dimensional Hamiltonian system is said to be *Liouville-integrable* if it admits n functionally independent first integrals in involution (that is their Poisson brackets commute) and such systems can be integrated by quadrature and enjoy a particularly simple topology (the flow lives on products of tori and cylinders). For systems of n first-order ordinary differential equations (ODEs) *algebraic integrability* is defined as the existence of $(n - 1)$ functionally independent first integrals that are algebraic functions of the dependent variables. The existence of such first integrals is important for integrability as they can be used to reduce the dimensionality of the system.

In the particular case of Hamiltonian systems with more than one degree of freedom, algebraic integrability corresponds to the notion of superintegrability (see Refs. 6, 7, and 16 and references therein). For a $2n$ -dimensional Hamiltonian system with $n > 1$, the $2n - 1$ first integrals needed for the property of algebraic integrability are more than the n ones needed for complete Liouville–Arnold integrability.

A different notion of integrability was introduced by Painlevé in the beginning of the 20th century.^{8,25} Although Painlevé's goal was not to define a notion of integrability but rather to build new functions, his property is today widely used for the detection of integrable systems. An ODE is said to possess the *Painlevé property*^{1,9,21} if its general solution is single-valued in its maximal domain of analytic continuation. The restrictions the Painlevé property impose on the solutions are so strong that an ODE exhibiting it may be considered for all practical purposes integrable. Many formal links between the Painlevé property and other notions of integrability have been established (see, for example, Refs. 9 and 10). Despite the fact that there is no general algorithmic way to obtain sufficient conditions for a given ODE to have the Painlevé property, necessary conditions can be derived. The algorithmic procedure to obtain such conditions is known as the *Painlevé test*

^{a)}Electronic mail: lafortus@math.arizona.edu

^{b)}Electronic mail: goriely@math.arizona.edu

or *singularity analysis* and is based on a local analysis of the solutions around movable isolated singularities.

The notions of algebraic integrability and singularity analysis of the solutions can be extended to finite difference equations. One of the most effective ways to perform singularity analysis in this context is given by *singularity confinement*^{5,10–15,17,27–29} which, like the Painlevé test for ODEs, imposes conditions on the singularities of the solutions. Although there exist many striking particular examples which indicate that singularity confinement is closely related to other notions of integrability, there is no clear-cut result which establishes this relationship in a formal and general framework. This article explores the relationship between the notion of algebraic integrability and the property of singularity confinement in discrete mappings.

In the case of systems of ODEs, Yoshida^{9,32,33} proved that the degree of a rational first integral is closely related to some exponents that can be obtained through the application of the Painlevé test. In this article, we establish an equivalent result for discrete systems by showing that singularity confinement analysis provides necessary conditions for the existence of an algebraic first integral. Moreover, a lower bound on the degree of rational first integrals can be obtained.

The rest of this article is organized as follows. In Sec. II, we define the notion of singularity and singularity confinement for birational autonomous discrete dynamical systems. In Sec. III, we study the local consequences of the existence of an algebraic first integral on the confinement of the singularities in the two-dimensional case. In Sec. IV, we extend the results of Sec. III to arbitrary dimensions. Applications are discussed in Sec. V.

II. FORMULATION OF THE PROBLEM

In this section we introduce and illustrate the notion of singularity confinement by considering a particular class of two-dimensional autonomous dynamical systems. These examples are used to motivate a formal definition valid in arbitrary dimensions.

Throughout the article the following notation will be used: If \mathbf{g} is an analytic function from \mathbb{C}^p to \mathbb{C}^r , then $D\mathbf{g}(\mathbf{x})$ is its *Jacobian* matrix evaluated at $\mathbf{x} \in \mathbb{C}^p$. When $p=r$, the *Jacobian* is denoted by $\det(D\mathbf{g}(\mathbf{x}))$. If f is a complex-valued rational function on \mathbb{C}^p , then $\text{num}(f)$ and $\text{den}(f)$ respectively denote the numerator and denominator of f .

A. Simple examples

The basic idea of singularity confinement is to consider the properties of solutions close to some singularities. To illustrate this concept, consider the class of two-dimensional complex dynamical systems of the form

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} g(x_n, y_n) \\ x_n \end{pmatrix}, \quad (2.1)$$

where g is a complex-valued rational function on \mathbb{C}^2 ,

$$g(x, y) = \frac{p(x, y)}{q(x, y)},$$

and p, q are relatively prime polynomials. Two types of singularities can be distinguished for such systems. The singularities of *first type* are the roots of q in \mathbb{C}^2 , that is, the values (x, y) at which the vector field $\mathbf{f}(x, y)$ in (2.1) has a singularity. The singularities of *second type* are the points where the Jacobian of \mathbf{f} is zero (this type of singularity was first considered in Ref. 12). As an example, consider the following discrete dynamical system:

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} -x_n - y_n + a + \frac{b}{x_n} \\ x_n \end{pmatrix}, \quad (2.2)$$

where a and b are complex numbers. The analysis is performed for generic values of the parameters a and b . Clearly the right-hand-side of (2.2) presents a singularity if $x_n=0$. To study this singularity we study the behavior of nearby solutions by introducing a small complex perturbation ϵ , with $|\epsilon|>0$. We then set y_0 to be any complex number and $x_0=\epsilon$. The iterates x_p and y_p are then determined for any positive nonzero integer p by the discrete dynamical system (2.2). After one iteration, we have

$$x_1 = \frac{b}{\epsilon} + a - y_0 + \mathcal{O}(\epsilon). \tag{2.3}$$

In the limit $\epsilon \rightarrow 0, |x_1| \rightarrow \infty$ and, using (2.2) again,

$$x_2 = -\frac{b}{\epsilon} + y_0 + \mathcal{O}(\epsilon), \tag{2.4}$$

$$x_3 = -\epsilon + \mathcal{O}(\epsilon^2). \tag{2.5}$$

The limit $\epsilon \rightarrow 0$ of the next iterate x_4 is well defined and given by y_0 . Despite the fact that the function \mathbf{f} defined by (2.2) is not well defined on any point of \mathbb{C}^2 of the form $(0,y)$, the limit $(x,y) \rightarrow (0,y)$ of $\mathbf{f}^4(x,y)$ exists and is given by

$$\lim_{(x,y) \rightarrow (0,y)} \mathbf{f}^4(x,y) = \begin{pmatrix} y \\ 0 \end{pmatrix}. \tag{2.6}$$

Moreover, since the value of the limit (2.6) depends on y , the same limit applied on the Jacobian of \mathbf{f}^4 is nonzero. The singularity is thus ‘‘confined’’ between the iterates 0 and 4 and does not propagate further. This property of confinement is very particular and we show in this article that it is closely related to the existence of a first integral for the system. A first integral for a discrete dynamical system of the form (2.1) is defined to be an analytic nonconstant complex-valued function I defined almost everywhere on \mathbb{C}^2 which is preserved by \mathbf{f} , that is

$$I(\mathbf{f}(\mathbf{x})) = I(\mathbf{x}), \tag{2.7}$$

for every $\mathbf{x} \in \mathbb{C}^2$ for which (2.7) makes sense. It is important to note that if \mathbf{f} is not defined at a point \mathbf{x}^* where I is well-defined, then the value of I at \mathbf{x}^* is still preserved under \mathbf{f} in the limit sense, i.e.,

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} I(\mathbf{f}(\mathbf{x})) = I(\mathbf{x}^*). \tag{2.8}$$

This is a consequence of the fact that \mathbf{f} and I are continuous and defined almost everywhere in \mathbb{C}^2 . In the particular case of system (2.2), one can check that the following polynomial,

$$I(x,y) = xy(x+y-a) - b(x+y), \tag{2.9}$$

is a first integral. Moreover, following (2.8), the first integral I is preserved in the limit sense through the iterations of the singularity of (2.2). That is,

$$-by = I(0,y) = \lim_{(x,y) \rightarrow (0,y)} I(\mathbf{f}^i(x,y)), \quad i = 1,2,3,4. \tag{2.10}$$

Note that most integrable systems studied in this article are particular cases of the so-called QRT-family of mappings.^{15,26}

The confinement property for the solutions of (2.1) does not hold in general. Although there is no formal result establishing how rare this confinement property might be, the fact that it does not hold for most systems of the form (2.1) has been well established in the literature.^{11,13,14,27-29}

The following system is an example where singularities are not confined:

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} -x_n - y_n + a + \frac{b}{x_n^3} \\ x_n \end{pmatrix}. \tag{2.11}$$

Consider the singularity $(x^*, y^*) = (0, y)$ and, as before, introduce $x_0 = \epsilon$ and $y_0 \in \mathbb{C}$ nonvanishing. The next iterates are $x_1 = b/\epsilon^3 + (a - y_0) - \epsilon$, $x_2 = -b/\epsilon^3 + y_0 + \mathcal{O}(\epsilon^5)$, and $x_3 = \epsilon + \mathcal{O}(\epsilon^5)$. Further, we find that, unlike the previous example, the appropriate cancellations allowing for confinement do not occur and the next iterates are also diverging at $\epsilon = 0$, $x_4 = 2b/\epsilon^3 + (a - y_0) - \epsilon + \mathcal{O}(\epsilon^5)$, and $x_5 = -2b/\epsilon^3 + y_0 + \mathcal{O}(\epsilon^5)$. In general, the sequence of limits $\epsilon \rightarrow 0$ of the $|x_p|$ for $p > 0$ repeats the formal triplet $(\infty, \infty, 0)$ indefinitely. This assertion is proven by induction based on the equality

$$\mathbf{f}^3 \begin{pmatrix} -nb/\epsilon^3 + y_0 + \mathcal{O}(\epsilon^5) \\ nb/\epsilon^3 + (a - y_0) - \epsilon + \mathcal{O}(\epsilon^5) \end{pmatrix} = \begin{pmatrix} -(n+1)b/\epsilon^3 + y_0 + \mathcal{O}(\epsilon^5) \\ (n+1)b/\epsilon^3 + (a - y_0) - \epsilon + \mathcal{O}(\epsilon^5) \end{pmatrix} \tag{2.12}$$

for any positive integer n . Therefore, the singularity is not confined. It is generally believed that systems which lack the confinement property will not be integrable. However, there is no definite result attached to this belief. We prove in the next sections that the analysis of the dynamics near the singularities can be used to conclude that (2.11) does not admit an algebraic first integral.

The following example illustrates the occurrence of the second type of singularity. Consider

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} \frac{x_n - 1}{y_n} + 2x_n \\ x_n \end{pmatrix}. \tag{2.13}$$

At first sight, the point $(x_n, 0)$ seems to be the only singularity. However, the Jacobian of \mathbf{f} in (2.13) vanishes whenever $x_n = 1$. If $x_0 = 1$ and $y_0 \neq 0$, the next iterate as determined by (2.13) is given by $x_1 = 2$. We then have $x_2 = 5$, $x_3 = 12$. It is not difficult to see that x_i grows with respect to i , and that all iterates of $(1, y_0)$ under (2.13) with $y_0 \neq 0$ are independent of y_0 . The singularity here only appears in the inverse of the Jacobian matrix and is not confined.

In the light of the previous examples, we can define the confinement property for singularities of discrete dynamical systems of the form (2.1). A general definition will be given in the next section. A *singularity* of the dynamical system (2.1) is defined to be any point (x^*, y^*) in \mathbb{C}^2 at which the right-hand-side of (2.1) is undefined or at which the Jacobian of \mathbf{f} is zero. A singularity (x^*, y^*) is said to be *confined* if there exists a positive integer N such that both limits

$$\lim_{(x,y) \rightarrow (x^*, y^*)} \mathbf{f}^N(x, y), \quad \lim_{(x,y) \rightarrow (x^*, y^*)} \det(\mathbf{Df}^N(x, y)) \tag{2.14}$$

exist and the second limit is nonzero. The smallest number N having this property is referred to as the *confinement number*, that is the number of steps necessary for confinement.

The next example illustrates the case when a singularity of the second type is confined,

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} \frac{(x_n - 1/a)(x_n - a)}{y_n(x_n - b)(x_n - 1/b)} \\ x_n \end{pmatrix}, \tag{2.15}$$

where a and b are two nonzero distinct complex numbers which are also both distinct from $1/a$ and $1/b$. The right-hand-side (RHS) of (2.15) presents a singularity of the second type if $x_n = a$ because the Jacobian of \mathbf{f} vanishes. If $x_0 = a$ and $y_0 \neq 0$, the next iterate as determined by (2.15) will be $x_1 = 0$; then $x_2 = 1/a$ and x_3 takes an indeterminate form $0/0$. Again, we introduce a small complex perturbation ϵ , $|\epsilon| > 0$, then set $x_0 = \epsilon$ to find that when $\epsilon \rightarrow 0$,

$$x_3 \rightarrow y_0 + a - \frac{1}{a}, \tag{2.16}$$

and we conclude that the singularity is confined. The dynamical system (2.15) has singularity of the first type if $x_n = b$. Applying the same procedure as before with a perturbation ϵ , one finds that the limit $\epsilon \rightarrow 0$ of $|x_1|$, x_2 and x_3 gives, respectively, ∞ , $1/b$ and $by_0/(b + y_0(1 - b^2))$. Thus, any singularity of the form (b, y_0) with $y_0 \neq b/(b^2 - 1)$ is confined in three steps. System (2.15) admits the following rational first integral:

$$I(x, y) = \left(b + \frac{1}{b}\right)(x + y) + \left(a + \frac{1}{a}\right)\left(\frac{1}{x} + \frac{1}{y}\right) - \frac{(1 + x^2)(1 + y^2)}{xy}. \tag{2.17}$$

The following example shows that the existence of a first integral does not imply that all singularities are confined. The system

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} x_n^2/y_n \\ x_n \end{pmatrix} \tag{2.18}$$

has the following first integral:

$$I(x, y) = \frac{xy}{x^2 + y^2}. \tag{2.19}$$

The RHS of (2.18) has a pole of order 1 at any point of \mathbb{C}^2 of the form $(x, 0)$ with $x \neq 0$. As before, if we study this singularity by perturbation we conclude that there is no confinement since the Laurent expansions of the iterations of (x, ϵ) under the discrete system (2.18) around $\epsilon = 0$ are all divergent. Although this singularity is not confined, one notices that any point of the form $(x, 0)$ with $x \neq 0$ does not have a unique preimage under (2.18). Indeed, the expression defining \mathbf{f}^{-1} given by

$$\mathbf{f}^{-1}(x, y) = \begin{pmatrix} y \\ y^2/x \end{pmatrix} \tag{2.20}$$

has a singularity of second type at $(x, 0)$ and all the points of this form are mapped to $(0, 0)$ under \mathbf{f}^{-1} . This example shows that if one wishes to establish a relationship between singularity confinement and first integrals, particular care should be given in distinguishing different singularities.

B. Preliminaries

The notion of singularity confinement will now be defined in a more general setting. We consider p -dimensional autonomous discrete dynamical systems described by birational mappings on \mathbb{C}^p . A *birational mapping* on \mathbb{C}^p is a \mathbb{C}^p -valued rational function which is one-to-one almost everywhere in \mathbb{C}^p and whose inverse, where it exists, is also represented by a rational function. The corresponding discrete dynamical systems are written

$$\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n), \quad \mathbf{x}_n \in \mathbb{C}^p, \quad n \in \mathbb{Z}, \tag{2.21}$$

where \mathbf{f} is a birational mapping on \mathbb{C}^p with no explicit dependence on n . The i th *iterate* of $\mathbf{x}_0 \in \mathbb{C}^p$ under (2.21) is said to *exist* if the following limit,

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}_0} \mathbf{f}^i(\mathbf{x}), \tag{2.22}$$

exists. The i th iterate is then defined to be that limit and is denoted $\mathbf{F}^i(\mathbf{x}_0)$.

Definition: A **singularity of type I** for the discrete dynamical system $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ is a point $\mathbf{x}^* \in \mathbb{C}^p$ at which \mathbf{f} is not defined. The set of singularity of type I associated with the mapping \mathbf{f} in \mathbb{C}^p is denoted $S_I(\mathbf{f})$.

Definition: A **singularity of type II** for the discrete dynamical system $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ is a point $\mathbf{x}^* \in \mathbb{C}^p$ such that $\det(\mathbf{Df}(\mathbf{x}^*)) = 0$. The set of singularity of type II in \mathbb{C}^p is denoted $S_{II}(\mathbf{f})$.

In the particular case of birational mappings, we have

$$S_I(\mathbf{f}) = \{\mathbf{x}^* \in \mathbb{C}^p \mid \text{den}(f_i)(\mathbf{x}^*) = 0, \text{ for some } i \in \{1, 2, \dots, p\}\}, \tag{2.23}$$

$$S_{II}(\mathbf{f}) = \{\mathbf{x}^* \in \mathbb{C}^p \mid \det(\mathbf{Df}(\mathbf{x}^*)) = 0\}. \tag{2.24}$$

We call the sets $S_I(\mathbf{f})$ and $S_{II}(\mathbf{f})$ the *singular sets of first and second type*, respectively. The *singular set* $S(\mathbf{f})$ is defined as the union of both sets:

$$S(\mathbf{f}) = S_I(\mathbf{f}) \cup S_{II}(\mathbf{f}). \tag{2.25}$$

For both types of singularity we can define the property of confinement.

Definition: Let $\mathbf{x}^* \in \mathbb{C}^p$ be a singularity of type I or II for the system $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$. The singularity is said to be **confined** if, for some $i_c \in \mathbb{N}^{>0}$, the iterate $\mathbf{F}^{i_c}(\mathbf{x}^*)$ exists and $\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \det(\mathbf{DF}^{i_c}(\mathbf{x})) \neq 0$.

The lowest such i_c is referred to as the *confinement number* that is the number of steps necessary for confinement. From now on, we assume that \mathbf{f} is a birational mapping. The following lemma proves to be useful.

Lemma 2.1: The image of $S_{II}(\mathbf{f})$ under the mapping \mathbf{f} lies inside an algebraic variety of codimension 2.

That is, the birational mapping \mathbf{f} is one-to-one only on the subset of \mathbb{C}^p defined by

$$O_{\mathbf{f}} \equiv \mathbb{C}^p \setminus S(\mathbf{f}). \tag{2.26}$$

As a consequence, the function \mathbf{f} restricted to $O_{\mathbf{f}}$ defines a rational diffeomorphism

$$\mathbf{f}: O_{\mathbf{f}} \rightarrow O'_{\mathbf{f}} \equiv \mathbf{f}(O_{\mathbf{f}}), \tag{2.27}$$

whose inverse is also a rational function.

Proof: Assume that $S_{II}(\mathbf{f}) \neq \emptyset$. Let $\mathbf{x}_0 \in S_{II}(\mathbf{f})$ and let $\mathbf{x}_1 = \mathbf{f}(\mathbf{x}_0)$. From the inverse function theorem, it follows that

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}_1} \|\det(\mathbf{Df}^{-1}(\mathbf{x}))\| = \infty. \tag{2.28}$$

Thus, for some i 's in $\{1, 2, \dots, p\}$, the denominator of $(\mathbf{f}^{-1})_i$ evaluated at \mathbf{x}_1 is zero. Denote those i by i_k , $k = 1, 2, \dots, r$. We now prove that at least for one i_k , $\text{num}((\mathbf{f}^{-1})_{i_k})$ also vanishes at \mathbf{x}_1 . By contradiction, suppose that none of these numerators vanishes at \mathbf{x}_1 . Thus

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}_1} \|\mathbf{f}^{-1}(\mathbf{x})\| = \infty. \tag{2.29}$$

On the other hand, we have

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}_0} \|\mathbf{f}^{-1}(\mathbf{f}(\mathbf{x}))\| = \mathbf{x}_0, \tag{2.30}$$

which contradicts Eq. (2.29). Thus, \mathbf{x}_1 must be in one of the following sets:

$$A_i = \{\mathbf{x} \in \mathbb{C}^p \mid \text{num}((\mathbf{f}^{-1})_i)(\mathbf{x}) = \text{den}((\mathbf{f}^{-1})_i)(\mathbf{x}) = 0\}, \quad i \in \{1, 2, \dots, p\}. \quad (2.31)$$

This concludes the proof since each set A_i is defined as the intersection of the zero level sets of two relatively prime polynomials. \square

As mentioned earlier, the singularity confinement turns out to be closely related to the existence of a first integral. This relationship will be explored in the next section.

Definition: A **first integral** of $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ is an analytic function $I: U \subset \mathbb{C}^p \rightarrow \mathbb{C}$ where U is a dense subset of \mathbb{C}^p such that

- (a) $\|(\partial I / \partial x_1, \partial I / \partial x_2, \dots, \partial I / \partial x_p)\| \neq 0$ almost everywhere in U , and
- (b) $I(\mathbf{f}(\mathbf{x})) = I(\mathbf{x})$.

A first integral can exist in several forms.

Definition: A first integral I of $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ is **polynomial** (resp. **rational**) if the function $I(\mathbf{x})$ is a polynomial (resp. rational) function of $\mathbf{x} \in \mathbb{C}^p$. A complex-valued function $f(\mathbf{x})$ is **algebraic** over \mathbb{C} if there exist $s > 0$ and q_0, \dots, q_s rational in \mathbf{x} such that

$$q_0 + q_1 f + \dots + q_s f^s = 0. \quad (2.32)$$

If s is the smallest positive integer such that (2.32) holds, the relation (2.32) is referred to as the minimal polynomial of f . A first integral I is **algebraic** if the function $I(\mathbf{x})$ is algebraic.

Many notions of integrability are used in the literature. In this article, we focus on algebraic integrability. We are interested in determining precisely “how many” algebraic first integrals a given discrete system must admit to be algebraically integrable. For instance, suppose that the algebraic functions $A_1(\mathbf{x}), A_2(\mathbf{x}), \dots, A_r(\mathbf{x})$ defined on \mathbb{C}^p are all first integrals for a given system and $F(z_1, z_2, \dots, z_r)$ is algebraic on \mathbb{C}^r . Then, it is clear that $A(\mathbf{x}) = F(A_1(\mathbf{x}), A_2(\mathbf{x}), \dots, A_r(\mathbf{x}))$ is also a first integral. However, this new first integral adds no knowledge to the given problem since it depends on the other first integrals.

Definition: Let $A_1(\mathbf{x}), A_2(\mathbf{x}), \dots, A_r(\mathbf{x})$ be smooth complex-valued functions defined on a domain $D \subset \mathbb{C}^p$. Then

- (a) A_1, A_2, \dots, A_r are **functionally dependent** if, for each $\mathbf{x} \in D$, there is a neighborhood U of \mathbf{x} and a smooth complex-valued function $F(z_1, z_2, \dots, z_r)$ not identically zero on any subset of \mathbb{C}^p such that

$$F(A_1(\mathbf{x}), A_2(\mathbf{x}), \dots, A_r(\mathbf{x})) = 0. \quad (2.33)$$

- (b) A_1, A_2, \dots, A_r are **functionally independent** if they are not functionally dependent when restricted to any open subset of D .

A simple way to determine if a set of functions is functionally independent is given by the following theorem (see, for example, Ref. 24).

Theorem: Let $\mathbf{A} = (A_1, A_2, \dots, A_r)$ be a smooth function from a domain $D \subseteq \mathbb{C}^p$ to \mathbb{C}^r . Then, $A_1(\mathbf{x}), A_2(\mathbf{x}), \dots, A_r(\mathbf{x})$ are functionally dependent if and only if $D\mathbf{A}(\mathbf{x}^*)$ has rank strictly less than r for all $\mathbf{x}^* \in D$.

We introduce the notion of integrability considered in this article.

Definition: The discrete dynamical system $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ is said to be **algebraically integrable** if it admits $(p - 1)$ algebraic functionally independent first integrals.

The following theorem is a direct extension of Brun’s^{9,23} theorem to discrete systems and is useful to extend the results we obtain for rational to algebraic first integrals.

Theorem: If a discrete dynamical system of $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ has k functionally independent algebraic first integrals, then it has k functionally independent rational first integrals.

Proof: Let I be an algebraic first integral and let

$$P(I) = q_0 + q_1 I + \cdots + q_{s-1} I^{s-1} + I^s \quad (2.34)$$

be its minimal polynomial, where q_i is a rational function of \mathbf{x} . Since I depends nontrivially on \mathbf{x} , there exists i such that q_i is a nonconstant rational function of \mathbf{x} . Since $I(\mathbf{f}(\mathbf{x})) = I(\mathbf{x})$, we have

$$q_0(\mathbf{f}(\mathbf{x})) - q_0(\mathbf{x}) + (q_1(\mathbf{f}(\mathbf{x})) - q_1(\mathbf{x}))I(\mathbf{x}) + \cdots + (q_{s-1}(\mathbf{f}(\mathbf{x})) - q_{s-1}(\mathbf{x}))I^{s-1}(\mathbf{x}) = 0. \quad (2.35)$$

Since P is minimal, we have $q_i(\mathbf{f}(\mathbf{x})) = q_i(\mathbf{x})$ and each q_i is a first integral. Now, let I' be another independent first integral whose minimal polynomial is

$$P(I') = q'_0 + q'_1 I' + \cdots + q'_{s'-1} I'^{s'-1} + I'^{s'}. \quad (2.36)$$

The independence between the two first integrals implies that there exist $i < s, i' < s'$ such that $q_i, q'_{i'}$ are two independent nonconstant rational first integrals. By induction, one can build k independent rational first integrals. \square

Therefore, algebraic integrability implies rational integrability and it is sufficient to consider discrete dynamical systems of the form (2.21) which admit $(p-1)$ functionally independent rational first integrals $R_i(\mathbf{x}) = P_i(\mathbf{x})/Q_i(\mathbf{x})$ where $i = 1, 2, \dots, p-1$ and P_i and Q_i are relatively prime polynomials. Let $L_{i,c}$ be the level set of the first integral R_i corresponding to the value $c \in \mathbb{C}$. Since \mathbf{f} is continuous, it leaves invariant the closure of the level set given by

$$\bar{L}_{i,c} = \{\mathbf{x} \in \mathbb{C}^p \mid P_i(\mathbf{x}) - c Q_i(\mathbf{x}) = 0\}. \quad (2.37)$$

III. THE TWO-DIMENSIONAL CASE

In this section we study the local implications of algebraic integrability on the confinement of singularities for two-dimensional discrete dynamical systems. We consider a discrete dynamical systems of the form (2.21) which is algebraically integrable, that is, it possesses a nontrivial rational first integral $R(\mathbf{x}) = P(\mathbf{x})/Q(\mathbf{x})$ where P and Q are some polynomials over \mathbb{C}^2 . Let L_c be the level set of R corresponding to the value $c \in \mathbb{C}$.

We first focus our attention on singularities of the first type. The singular set of first type $S_I(\mathbf{f})$ is an algebraic variety since it is defined as the set of zeros of polynomials. We consider its irreducible decomposition is

$$S_I(\mathbf{f}) = \bigcup_{i=1}^d S_I^{(i)}(\mathbf{f}), \quad (3.1)$$

where each irreducible component $S_I^{(i)}(\mathbf{f})$ is defined as the zero set of an irreducible polynomial on \mathbb{C}^2 .

In example (2.18), we showed that the existence of a rational first integral does not ensure that all singularities are confined. However, in this case it is not possible to “enter” the singularity, meaning that it does not have a unique preimage under the mapping. Moreover, we know that confinement is a generic property, that is, it is only concerned with dense subsets of irreducible components of S_I . For instance, in example (2.15) all singularities of type I of the form (b, y_0) confine in three steps except a particular singularity defined by $y_0 = b/(b^2 - 1)$. The next theorem relates the confinement of singularities to the existence of a rational first integral.

Theorem 3.1: Consider a two-dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and assume it has a rational first integral $R(\mathbf{x}) = P(\mathbf{x})/Q(\mathbf{x})$. Then, for each irreducible component $S_I^{(j)}(\mathbf{f})$, we have either

(a) *there exists $k_j \in \mathbb{N}^{>0}$ such that*

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \mathbf{f}^{k_j}(\mathbf{x}) \tag{3.2}$$

exists for almost all $\mathbf{x}^ \in S_I^{(j)}(\mathbf{f})$, or*

(b) *there exists $l_j \in \mathbb{N}^{>0}$ such that $S_I^{(j)}(\mathbf{f})$ lies inside the singular set of the birational mapping defining \mathbf{f}^{-l_j} .*

Proof: Without loss of generality, take $j = 1$. Consider two cases:

(1) $S_I^{(1)}(\mathbf{f}) \not\subset \bar{L}_c$ for all $c \in \mathbb{C}$.

Let $\mathbf{x}^* \in S_I^{(1)}(\mathbf{f})$ such that $\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \|f_i(\mathbf{x})\|$ exists or is infinity (as opposed to undefined) for $i = 1, 2$. Suppose that $\mathbf{x}^* \in \bar{L}_{c^*}$ for only one $c^* \in \mathbb{C}$ and that there is a neighborhood of \mathbf{x}^* on \bar{L}_{c^*} whose only intersection with any of the singular sets associated with the mappings \mathbf{f}^i , $i = 1, 2, \dots, q + 1$, (q is defined below) is \mathbf{x}^* itself. The set of points satisfying the above properties is dense in $S_I^{(1)}(\mathbf{f})$. Let $\mathbf{p}_i(\epsilon)$, $i = 1, 2, \dots, q$, be the paths in \mathbb{C}^2 such that

(i) $\mathbf{p}_i(\epsilon) \in \bar{L}_{c^*}$ for ϵ small enough, that is,

$$P(\mathbf{p}_i(\epsilon)) - c^* Q(\mathbf{p}_i(\epsilon)) = 0; \text{ and} \tag{3.3}$$

(ii) $\|\mathbf{p}_i(\epsilon)\| \rightarrow \infty$ as $\epsilon \rightarrow 0$ and at least one of the components of $\mathbf{p}_i(\epsilon)$ is of the form $1/\epsilon$.

Note that the number of paths q can depend on the complex number c^* but the values it can take for different c^* can be bounded above by a number depending only on the degrees of P and Q .

Now, consider another path $\mathbf{x}(\sigma) \in \bar{L}_{c^*} \forall \sigma$ such that

$$\lim_{\sigma \rightarrow 0} \mathbf{x}(\sigma) = \mathbf{x}^*. \tag{3.4}$$

Since there is a neighborhood of \mathbf{x}^* on \bar{L}_{c^*} whose only intersection with $S(\mathbf{f})$ is \mathbf{x}^* , $\mathbf{f}(\mathbf{x}(\sigma))$ is well-defined for σ nonzero and small enough. Moreover, since \mathbf{f} preserves \bar{L}_{c^*} , $\mathbf{f}(\mathbf{x}(\sigma))$ also lies in \bar{L}_{c^*} , and

$$\lim_{\sigma \rightarrow 0} \|\mathbf{f}(\mathbf{x}(\sigma))\| = \infty. \tag{3.5}$$

Therefore, there exists a change of variable $\sigma = \sigma(\epsilon)$ of order $\mathcal{O}(\epsilon^s)$ for some $s > 0$, so that $\mathbf{f}(\mathbf{x}(\sigma))$ is one of the paths \mathbf{p}_i defined above:

$$\mathbf{f}(\mathbf{x}(\sigma(\epsilon))) = \mathbf{p}_i(\epsilon), \tag{3.6}$$

for some $1 \leq i \leq q$. Our claim is that there exists m with $m \leq q + 1$ such that the limit as $\sigma \rightarrow 0$ of $\mathbf{f}^m(\mathbf{x}(\sigma))$ exists and is finite. Suppose, by contradiction, that the iterates $\mathbf{f}^k(\mathbf{x}(\sigma))$, $k = 1, 2, 3, \dots, q + 1$, are all divergent at $\sigma = 0$. Each iterate $\mathbf{f}^k(\mathbf{x}(\sigma))$ is associated with a unique path $\mathbf{p}_{i_k}(\epsilon)$. This implies that the k th preimage of $\mathbf{p}_{i_k}(\epsilon)$ is finite in the limit $\epsilon \rightarrow 0$ and no other $\mathbf{f}^l(\mathbf{x}(\sigma))$ for $1 \leq k < l \leq q + 1$ corresponds to the same $\mathbf{p}_{i_k}(\epsilon)$. Hence the relation between the $q + 1$ iterates that are divergent at $\sigma = 0$ and the paths $\mathbf{p}_i(\epsilon)$ is injective. This is a contradiction since there are only q paths $\mathbf{p}_i(\epsilon)$.

(2) There exists $c \in \mathbb{C}$ such that $S_I^{(1)}(\mathbf{f}) \subset \bar{L}_c$. Let

$$\bar{L}_c = \bigcup_{i=1}^r \bar{L}_c^{(i)} \tag{3.7}$$

be the irreducible decomposition of \bar{L}_c for any $c \in \mathbb{C}$. If $S_I^{(1)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-1} , then the theorem is proved. Otherwise, let $k \in \mathbb{N}^{>0}$ be such that $S_I^{(1)}(\mathbf{f})$ does not lie in a singular set associated with the mappings \mathbf{f}^{-m} , $m = 1, 2, \dots, k$. Then,

$$\overline{\mathbf{f}^{-m}(S_l^{(1)}(\mathbf{f}))} = \bigcup_{i \in B_m} \overline{L_c^{(i)}}, \quad 1 \leq l \leq k, \tag{3.8}$$

where each B_m is a subset of $\{1, 2, \dots, r\}$. The k sets in (3.8) are distinct, otherwise, by contradiction, there exist $m_1, m_2 \in \mathbb{N}^{>0}$ such that

$$\overline{\mathbf{f}^{-m_1}(S_l^{(1)}(\mathbf{f}))} = \overline{\mathbf{f}^{-m_2}(S_l^{(1)}(\mathbf{f}))}, \quad 1 \leq m_1 < m_2 \leq k. \tag{3.9}$$

This implies that

$$S_l^{(1)}(\mathbf{f}) = \overline{\mathbf{f}^{-1}(\mathbf{f}^{m_1 - m_2 + 1}(S_l^{(1)}(\mathbf{f})))}, \tag{3.10}$$

which is a contradiction since \mathbf{f} is undefined almost everywhere on $S_l^{(1)}(\mathbf{f})$. Hence the k sets in (3.8) are distinct. However, there are only finitely many possibilities for the sets B_m in (3.8). Hence $S_l^{(1)}(\mathbf{f})$ must lie inside the singular set of \mathbf{f}^{-l_1} for some l_1 . \square

Notice that condition (3.2) does not imply that the singularity is confined since the condition that the Jacobian be nonzero must be satisfied. In order to solve this problem, singularities of type II have to be considered. The singular set of second type is not, in general, an algebraic variety but its closure,

$$\overline{S_{II}(\mathbf{f})} = \{\mathbf{x}^* \in \mathbb{C}^p \mid \text{num}(\det(D\mathbf{f}(\mathbf{x}^*))) = 0\}, \tag{3.11}$$

is. Hence, it is possible to introduce the irreducible decomposition

$$\overline{S_{II}(\mathbf{f})} = \bigcup_{i=1}^s \overline{S_{II}^{(i)}(\mathbf{f})}, \tag{3.12}$$

where each $S_{II}^{(i)}(\mathbf{f})$ is a subset of $S_{II}(\mathbf{f})$. It is now possible to state the general theorem for singularities of second type.

Theorem 3.2: Consider a two-dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and assume it has a rational first integral $R(\mathbf{x}) = P(\mathbf{x})/Q(\mathbf{x})$. Then, for each irreducible component $\overline{S_{II}^{(j)}(\mathbf{f})}$, we have either

(a) there exists $k_j \in \mathbb{N}^{>0}$ such that

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \det(D\mathbf{f}^{k_j}(\mathbf{x})) \tag{3.13}$$

exists and is nonzero for almost all $\mathbf{x} \in S_{II}^{(j)}(\mathbf{f})$, or

(b) there exists $l_j \in \mathbb{N}^{>0}$ such that $S_{II}^{(j)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-l_j} .

Proof: The proof is similar to the proof of Theorem 3.1. Again, without loss of generality, we take $j = 1$ and consider two cases.

(1) $\overline{S_{II}^{(1)}(\mathbf{f})} \not\subset \overline{L_c}$ for all $c \in \mathbb{C}$.

The image of $S_{II}^{(1)}(\mathbf{f})$ under the mapping \mathbf{f} is a finite subset of \mathbb{C}^2 , thanks to Lemma 2.1. Moreover, $S_{II}^{(1)}(\mathbf{f})$ is not contained inside any level set. For any point in the image of $S_{II}^{(1)}(\mathbf{f})$ it is therefore possible to choose two distinct preimages in $S_{II}^{(1)}(\mathbf{f})$ which lie in two different level sets. Since \mathbf{f} preserves $\overline{L_c}$, every point in the image of $S_{II}^{(1)}(\mathbf{f})$ must lie in the set

$$\bigcap_{c \in \mathbb{C}} \overline{L_c} = \{\mathbf{x} \in \mathbb{C}^2 \mid P(\mathbf{x}) = Q(\mathbf{x}) = 0\}. \tag{3.14}$$

The elements of this finite set are denoted \mathbf{s}_i , $i = 1, 2, \dots, m$. Note that in the case of a polynomial first integral, $Q(\mathbf{x}) = 1$ and this set is empty.

Now, choose $\mathbf{x}^* \in S_I^{(1)}(\mathbf{f})$ such that $\mathbf{x}^* \in \bar{L}_{c^*}$ for only one $c^* \in \mathbb{C}$ and there is a neighborhood of \mathbf{x}^* on \bar{L}_{c^*} whose only intersection with any of the singular sets associated with the mappings \mathbf{f}^i , $i = 1, 2, \dots, q+r+1$, (q and r are defined below) is \mathbf{x}^* itself. The set of points satisfying the above properties is dense in $S_I^{(1)}(\mathbf{f})$. Consider the q paths $\mathbf{p}_i(\epsilon)$ described in the proof of Theorem 3.1. Consider also the r paths $\mathbf{w}_i(\epsilon)$, $i = 1, 2, \dots, r$, in \mathbb{C}^2 satisfying

- (i) $\mathbf{w}_i(\epsilon) \in \bar{L}_{c^*}$ for ϵ small enough.
- (ii) $\mathbf{w}_i(\epsilon) \rightarrow \mathbf{s}_{j_i}$ as $\epsilon \rightarrow 0$ and one of the components ($k = 1$ or 2) of $\mathbf{w}_i(\epsilon)$ is of the form $(\mathbf{s}_{j_i})_k + \epsilon$ for some $1 \leq j_i \leq m$.

Now, consider another path $\mathbf{x}(\sigma) \in \bar{L}_{c^*}$ such that

$$\lim_{\sigma \rightarrow 0} \mathbf{x}(\sigma) = \mathbf{x}^*. \tag{3.15}$$

Since there is a neighborhood of \mathbf{x}^* on \bar{L}_{c^*} whose only intersection with $S(\mathbf{f})$ is \mathbf{x}^* , $\mathbf{f}(\mathbf{x}(\sigma))$ is well-defined for σ nonzero and small enough. Moreover, since \mathbf{f} preserves \bar{L}_{c^*} , $\mathbf{f}(\mathbf{x}(\sigma)) \in \bar{L}_{c^*}$, and

$$\lim_{\sigma \rightarrow 0} \mathbf{f}(\mathbf{x}(\sigma)) = \mathbf{s}_i \tag{3.16}$$

for some i such that $1 \leq i \leq r$. Following the proof of Theorem 3.1, $\mathbf{f}(\mathbf{x}(\sigma))$ can be associated with a path $\mathbf{w}_j(\epsilon)$ (whose limit as $\sigma \rightarrow 0$ is \mathbf{s}_i). Suppose by contradiction that the $d+r+1$ iterates $\mathbf{f}^k(\mathbf{x}(\sigma))$, $k = 1, 2, \dots, d+r+1$, are either divergent at $\sigma=0$ or have a limit $\sigma \rightarrow 0$ equal to one of the \mathbf{s}_i . Each iterate \mathbf{f}^k is associated with a path $\mathbf{p}_{i_k}(\epsilon)$ or $\mathbf{w}_{i_k}(\epsilon)$. This implies that the unique k th preimage of $\mathbf{p}_{i_k}(\epsilon)$ or $\mathbf{w}_{i_k}(\epsilon)$ must converge to \mathbf{x}^* as $\epsilon \rightarrow 0$. The preimage is unique because there is a neighborhood of \mathbf{x}^* on \bar{L}_{c^*} whose only intersection with any of the singular sets associated with the mappings \mathbf{f}^i , $i = 1, 2, \dots, q+r+1$, (q and r are defined below) is \mathbf{x}^* itself. Since \mathbf{x}^* is not one of the \mathbf{s}_i , there cannot be two iterates $\mathbf{f}^{k_1}(\mathbf{x}(\sigma))$ and $\mathbf{f}^{k_2}(\mathbf{x}(\sigma))$, $k_1, k_2 \leq q+r+1$, associated with the same path. Since there are only $q+r+1$ paths $\mathbf{p}_i(\epsilon)$ and $\mathbf{w}_i(\epsilon)$, there exists a $k \leq q+r+1$ such that the limit $\sigma \rightarrow 0$ of the k th iterate of $\mathbf{x}(\sigma)$ must be finite and not equal to one \mathbf{s}_i as $\sigma \rightarrow 0$. Therefore, the limit $\sigma \rightarrow 0$ of the Jacobian of \mathbf{f}^k evaluated at $\mathbf{x}(\sigma)$ must be a finite nonvanishing number. This ends the proof in the first case.

(2) There exists $c \in \mathbb{C}$ such that $S_I^{(1)}(\mathbf{f}) \subset \bar{L}_c$. The argument leading to the conclusion that $S_I^{(j)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-l_1} for some l_1 is identical to the second case considered in the proof of the preceding theorem. \square

With Theorem 3.2, it is now easy to see that singularities of type I satisfying the condition (3.2) are confined. Indeed, suppose k_j is the number considered in the first part of Theorem 3.1. If the Jacobian of \mathbf{f}^{k_j} is zero almost everywhere on $S_I^{(j)}(\mathbf{f})$, then $S_I^{(j)}(\mathbf{f})$ becomes a subset of $\overline{S_{II}(\mathbf{f}^{k_j})}$. According to the proof of Theorem 3.1 we know that $S_I^{(j)}(\mathbf{f}) \not\subset \bar{L}_c$ for all $c \in \mathbb{C}$. Thus, as a consequence of Theorem 3.2, the singularities in $S_I^{(j)}(\mathbf{f})$ must be confined. We can state a general theorem including both types of singularities. To do so, we introduce the irreducible decomposition associated with $S(\mathbf{f})$

$$S(\mathbf{f}) = \bigcup_{i=1}^{d+s} S^{(i)}(\mathbf{f}), \tag{3.17}$$

where d and s are defined in (3.1) and (3.12).

Theorem 3.3: Consider a two-dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and assume it has a rational first integral $R(\mathbf{x}) = P(\mathbf{x})/Q(\mathbf{x})$. Let $S(\mathbf{f})$ be its singular set with irreducible components $S^{(j)}(\mathbf{f})$. Then, for each j , either

- (a) almost all singularities of $S^{(j)}(\mathbf{f})$ are confined in the same number of steps, or
- (b) there exists $l_j \in \mathbb{N}^{>0}$ such that $S^{(j)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-l_j} .

IV. ARBITRARY DIMENSIONS

In this section, we extend the results of the previous section to birational mappings of arbitrary dimension p . The mappings are assumed to be algebraically integrable, that is, they admit $(p - 1)$ functionally independent first integrals $R_i(\mathbf{x}) = P_i(\mathbf{x})/Q_i(\mathbf{x})$, $i = 1, 2, \dots, p - 1$. Then, the closure of the level sets associated with $\mathbf{c} \in \mathbb{C}^{p-1}$ is given by

$$\bar{L}_{\mathbf{c}} = \{\mathbf{x} \in \mathbb{C}^p \mid P_i(\mathbf{x}) - c_i Q_i(\mathbf{x}) = 0, \text{ for } 1 \leq i \leq p - 1\} = \bigcap_{i=1}^{p-1} \bar{L}_{i,c_i}, \tag{4.1}$$

with \bar{L}_{i,c_i} defined in (2.37). A set of constants $\mathbf{c} \in \mathbb{C}^{p-1}$ is said to be *regular* for the first integrals $\mathbf{R} = (R_1, R_2, \dots, R_{p-1})$ if $D\mathbf{R}(\mathbf{x})$ has rank equal to $(p - 1)$ almost everywhere on $\mathbf{R}^{-1}(\mathbf{c})$. The set $\bar{L}_{\mathbf{c}}$ is then an algebraic variety of codimension $(p - 1)$.

Our main Theorems 3.1–3.3 can be readily generalized to arbitrary dimensions and we only give here an outline of the proofs.

Theorem 4.1: Consider a p -dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and assume it is algebraically integrable with first integrals $R_i(\mathbf{x}) = P_i(\mathbf{x})/Q_i(\mathbf{x})$, $i = 1, 2, \dots, p - 1$. Then for each irreducible component $S_I^{(j)}(\mathbf{f})$, we have either

- (a) there exists $k_j \in \mathbb{N}^{>0}$ such that

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \mathbf{f}^{k_j}(\mathbf{x}) \tag{4.2}$$

exists for almost all $\mathbf{x} \in S_I^{(j)}(\mathbf{f})$, or

- (b) there exists $l_j \in \mathbb{N}^{>0}$ such that $S_I^{(j)}(\mathbf{f})$ lies inside the singular set of the birational mapping defining \mathbf{f}^{-l_j} .

Proof: Without loss of generality, take $j = 1$. Consider two cases:

- (1) $S_I^{(1)}(\mathbf{f}) \not\subset \bar{L}_{i,c}$ for all $c \in \mathbb{C}$ and $1 \leq i \leq p - 1$.

Let $\mathbf{x}^* \in S_I^{(1)}(\mathbf{f})$ such that $\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \|f_i(\mathbf{x})\|$ exists or is infinite (as opposed to undefined) for $i = 1, \dots, p$. Suppose that $\mathbf{x}^* \in \bar{L}_{\mathbf{c}^*}$ for only one $\mathbf{c}^* \in \mathbb{C}^{p-1}$ which is regular and that there is a neighborhood of \mathbf{x}^* on $\bar{L}_{\mathbf{c}^*}$ whose only intersection with the singular sets associated with the mappings \mathbf{f}^i , $i = 1, 2, \dots, q + 1$, (q is defined below) is \mathbf{x}^* itself. The set of points satisfying the above properties is dense in $S_I^{(1)}(\mathbf{f})$. Let q be the number of paths $\mathbf{p}_i(\epsilon)$, $i = 1, 2, \dots, q$, in \mathbb{C}^p such that

- (i) $\mathbf{p}_i(\epsilon) \in \bar{L}_{\mathbf{c}^*}$ for ϵ small enough. That is,

$$P_k(\mathbf{p}_i(\epsilon)) - c_k^* Q_k(\mathbf{p}_i(\epsilon)) = 0, \quad k = 1, 2, \dots, p - 1. \tag{4.3}$$

- (ii) $\|\mathbf{p}_i(\epsilon)\| \rightarrow \infty$ as $\epsilon \rightarrow 0$ and at least one of the components of $\mathbf{p}_i(\epsilon)$ is of the form $1/\epsilon$.

Using the fact that $\bar{L}_{\mathbf{c}^*}$ is of codimension $p - 1$, the rest of the proof follows exactly the similar case in the proof of Theorem 3.1.

- (2) There exists $c \in \mathbb{C}$ and i , $1 \leq i \leq p - 1$, such that $S_I^{(1)}(\mathbf{f}) \subset \bar{L}_{i,c}$. This part of the proof is identical to the corresponding part in the proof of Theorem 3.1 except that $\bar{L}_{i,c}$ is considered instead of $\bar{L}_{\mathbf{c}}$. □

Theorem 4.2: Consider a p -dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and assume it is algebraically integrable with rational first integrals $R_i(\mathbf{x}) = P_i(\mathbf{x})/Q_i(\mathbf{x})$, $i = 1, 2, \dots, p-1$. Then, for each irreducible component $S_H^{(j)}(\mathbf{f})$, we have either

(a) there exists $k_j \in \mathbb{N}^{>0}$ such that

$$\lim_{\mathbf{x} \rightarrow \mathbf{x}^*} \det(D\mathbf{f}^{k_j}(\mathbf{x})) \tag{4.4}$$

exists and is nonzero for almost all $\mathbf{x} \in S_H^{(j)}(\mathbf{f})$, or

(b) there exists $l_j \in \mathbb{N}^{>0}$ such that $S_H^{(j)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-l_j} .

Proof: The proof is similar to the proof of Theorem 3.2 and the only difference comes in the first case, when $S_H^{(1)}(\mathbf{f}) \not\subset \bar{L}_{i,c}$ for all $c \in \mathbb{C}$ and $1 \leq i \leq p-1$. Following the proof of Theorem 3.2, for any point in the image of $S_H^{(1)}(\mathbf{f})$ it is possible to choose two distinct preimages in $S_H^{(1)}(\mathbf{f})$ which lie in two different level sets. Indeed, from Lemma 2.1, the image of $S_H^{(1)}(\mathbf{f})$ under the mapping \mathbf{f} lies in a subset of codimension 2 inside \mathbb{C}^p . Since \mathbf{f} preserves $\bar{L}_{i,c}$, the points in the image of $S_H^{(1)}(\mathbf{f})$ must lie in the set

$$\bigcap_{c \in \mathbb{C}, i \leq p-1} \bar{L}_{i,c} = \{\mathbf{x} \in \mathbb{C}^2 \mid P_i(\mathbf{x}) = Q_i(\mathbf{x}) = 0\}. \tag{4.5}$$

Since the first integrals are functionally independent, the set defined above is finite. The rest of the proof is similar to the proof of Theorem 3.2 except that one should specify that \mathbf{c}^* has to be regular. \square

Theorem 4.3: Consider a p -dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and assume it is algebraically integrable. Let $S(\mathbf{f})$ be its singular set with irreducible components $S^{(j)}(\mathbf{f})$. Then, for each j , either

- (a) almost all singularities of $S^{(j)}(\mathbf{f})$ are confined in the same number of steps, or
- (b) there exists $l_j \in \mathbb{N}^{>0}$ such that $S^{(j)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-l_j} .

V. COROLLARIES AND APPLICATIONS

In the previous two sections, the existence of first integrals was assumed to obtain local information on the confinement property. Here, the information given by the singularity confinement property is used to obtain global information on the discrete dynamical systems such as the nonexistence of algebraic first integral and the degree of possible rational first integrals. The results of this section can be considered as a discrete analog of Yoshida’s theorem^{9,32,33} which, for ODEs, relates the Kovalevskaya exponents given by the Painlevé test to the degree of a rational first integral. The first corollary is a direct consequence of Theorem 4.3.

Corollary 5.1: Consider a p -dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ with a nonempty singular set $S(\mathbf{f})$. If there exists an irreducible components $S^{(j)}(\mathbf{f})$ in which almost all singularities are not confining and such that $S^{(j)}(\mathbf{f}) \not\subset S(\mathbf{f}^{-k})$ for any positive k , then the system is not algebraically integrable.

As an example of this corollary, we show that system (2.11), defined by

$$\mathbf{f}(x,y) = \begin{pmatrix} -x - y + a + \frac{b}{x^3} \\ x \end{pmatrix}, \tag{5.1}$$

is not algebraically integrable. Recall that singularities of the form $(0,y)$ are nonconfining. In order to use Corollary 5.1 and prove that this mapping does not admit an algebraic first integral, one must show that points of the form $(0,y)$ are not generically in the singular set of \mathbf{f}^{-l} for some

$l > 0$. The singular set of \mathbf{f}^{-1} consists of points of the form $(x, 0)$. Thus, we must prove that generic iterates of $(0, y)$ under \mathbf{f}^{-1} do not belong to the set of points of the form $(x, 0)$. The inverse of the mapping (5.1) is given by

$$\mathbf{f}^{-1}(x, y) = \begin{pmatrix} y \\ -x - y + a + \frac{b}{y^3} \end{pmatrix}. \tag{5.2}$$

This mapping can be obtained from \mathbf{f} of (5.1) by interchanging x and y . Therefore, we can follow the analysis performed in (2.12) to perform a Laurent expansion in y to show that

$$\mathbf{f}^{-2}(x_0, y_0) = \begin{pmatrix} b/y_0^3 + (a - x_0) - y_0 + \mathcal{O}(y_0^5) \\ -b/y_0^3 + x_0 + \mathcal{O}(y_0^5) \end{pmatrix}. \tag{5.3}$$

Moreover, we have

$$\mathbf{f}^{-3} \begin{pmatrix} nb/y_0^3 + (a - x_0) - y_0 + \mathcal{O}(y_0^5) \\ -nb/y_0^3 + x_0 + \mathcal{O}(y_0^5) \end{pmatrix} = \begin{pmatrix} (n+1)b/y_0^3 + (a - x_0) - y_0 + \mathcal{O}(y_0^5) \\ -(n+1)b/y_0^3 + x_0 + \mathcal{O}(y_0^5) \end{pmatrix}, \tag{5.4}$$

where n is any positive integer. Hence, points of the form $(0, y)$ are not generically sent to points of the form $(x, 0)$ by applications of \mathbf{f}^{-1} and, from Corollary 5.1, system (2.11) does not admit an algebraic first integral.

In general, algebraic integrability restricts the possible *local* behavior of singularities. Theorem 4.3 gives two possible local behaviors but, in each case, the singularity can have several different *global* behaviors. Consider an irreducible component of the singular set $S^{(1)}(\mathbf{f})$. In the proofs of Theorems 3.3 and 4.3, we considered two cases: $S^{(1)}(\mathbf{f}) \not\subset \bar{L}_c$ and $S^{(1)}(\mathbf{f}) \subset \bar{L}_c$. When $S^{(1)}(\mathbf{f}) \not\subset \bar{L}_c$ the singularities of $S_I^{(1)}(\mathbf{f})$ are generically confined. In this case, two different types of global behavior can be expected. First, after meeting a finite number singularities, the mapping generically never meets other singularities (note that this must also be true in the backward direction, that is for \mathbf{f}^{-1}). We will refer to such a behavior as *global confinement*. Second, the singularities may be confined but not globally. Because, for each dynamical system of the form (2.21), there are only finitely many irreducible parts to $S(\mathbf{f})$, the mapping must, at one point, come back to the initial singularity and take $S_I^{(1)}(\mathbf{f})$ into itself. We say that such singularities are *periodic*. We now define formally these two types of global behavior and illustrate them.

Definition: A singularity $\mathbf{x}^* \in \mathbb{C}^p$ for a birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ is said to be **globally confined** if there exist $k', k'' \in \mathbb{N}^{>0}$ such that $\mathbf{x}^* \notin S(\mathbf{f}^k)$ for any $k > k'$ and $\mathbf{x}^* \in S(\mathbf{f}^{-k})$ for any $k > k''$.

An example of such a behavior is given by system (2.2). For this system, points of the form $(x, 0)$ are not sent to points of the form $(0, y)$ through successive applications of \mathbf{f} . To prove this, let (x_n, y_n) be the n th iterate of $(x, 0)$. If $n = 4k$, an expansion of $\mathbf{f}^4(x, 0)$ reveals that $(x_{4k}, y_{4k}) = (kx + \mathcal{O}(x^2), (-k+1)x + \mathcal{O}(x^2))$, $k = 1, 2, \dots$. This fact also implies that $x_{4k+3} = -kx + \mathcal{O}(x^2)$ and we conclude that x_{4k+3} and x_{4k} are neither infinite nor vanishing. By contradiction assume that there exists n such that $(x_n, y_n) = (0, y)$. Therefore, either $n = 4k + 1$ or $n = 4k + 2$. However, from Eq. (2.4) we know that x_{n+2} is infinite which is a contradiction. We conclude that no iterate of $(x, 0)$ falls onto a point of the form $(0, y)$ and singularities of the form $(0, y)$ are globally confined.

Definition: Consider the singular set of a birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ and one of its irreducible component $S^{(j)}(\mathbf{f})$. If there is a $k \in \mathbb{N}^{>0}$ such that $\mathbf{f}^k(S^{(j)}(\mathbf{f})) = S^{(j)}(\mathbf{f})$, then the elements of $S^{(j)}(\mathbf{f})$ are said to be **periodic singularities** of period k .

An example of such a behavior is given by a particular case of the Gambier system^{13,14}

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} (y_n + (1/a - a - 1))x_n - a + 1 \\ x_n + y_n \\ 1 - 1/y_n \end{pmatrix}, \tag{5.5}$$

where a is a nonzero complex constant. The system (5.5) admits a singularity of type II at any point of the form (x, a) . Then the first iterate of a point of this form is given by $(1/a - 1, 1 - 1/a)$. But the Jacobians of \mathbf{f}^2 and \mathbf{f}^3 at (x, a) are nonzero and the singularity is confined. It is not globally confined since $\mathbf{f}^4(x, a) = \mathbf{f}(x, a) = (1/a - 1, 1 - 1/a)$ and the singularity is periodic. The system admits the following first integral:

$$I(x, y) = \frac{y^3 - 3y + 1}{y(y - 1)}. \tag{5.6}$$

Now, consider the second case considered in the proofs of Theorems 3.3 and 4.3: $S^{(1)}(\mathbf{f}) \subset \bar{L}_{i,c}$ for some c . If the singularities of $S^{(1)}(\mathbf{f})$ are generically confined in k steps, then $\mathbf{f}^k(S^{(1)}(\mathbf{f}))$ has to be a set of the form of the RHS of (3.8). Since there are only finitely many of those, the singularities cannot be globally confined. So, either they are periodic, or $S^{(1)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^k for all k greater than a certain k' . It must also be true in the backward direction and if the singularities are not periodic $S^{(1)}(\mathbf{f})$ lies inside the singular set of \mathbf{f}^{-k} for all k greater than a certain k'' . We will refer to these singularities as *ubiquitous singularities*. An example of such a behavior is given by the singularities of Eq. (2.18) of the form $(x, 0)$. These singularities are not confined but they are in the singular set of \mathbf{f}^{-l} for any $l > 0$ and, therefore, these singularities are ubiquitous. Note that singularities of the form $(0, y)$ are also ubiquitous.

Definition: A singularity $\mathbf{x}^* \in \mathbb{C}^p$ of a birational mapping is said to be **ubiquitous** if there exist a $k' \in \mathbb{N}^{>0}$ such that $\mathbf{x}^* \in S(\mathbf{f}^k)$ for any $k > k'$ and a $k'' \in \mathbb{N}^{>0}$ such that $\mathbf{x}^* \in S(\mathbf{f}^{-k})$ for any $k > k''$.

Given a discrete dynamical system of the form (2.21) and an irreducible component $S^{(j)}(\mathbf{f})$ of the singular set, one would like to be able to obtain information on the first integrals. The only way one can know if a given irreducible component of the singular set lies inside a level set is by studying the global behavior of the singularity. If the singularity is globally confined, then $S^{(j)}(\mathbf{f})$ does not lie inside a level set, but if it is ubiquitous, then it does. If the singularities are periodic, no conclusion can be reached.

Information on first integrals can be obtained by studying singularities of type I. Consider the paths \mathbf{p}_i defined in the two-dimensional case in the proof of Theorem 3.1. The number of these paths is bounded above by the maximum of the degrees of P and Q . Moreover, for each $S_I^{(j)}(\mathbf{f}) \not\subset \bar{L}_c$ for all c , consider k'_j , the lowest value k_j can take. Then $k'_j - 1$ is the number of different paths \mathbf{p}_i realized by iterating a point close to the singularity under the mapping. Hence, the following corollary holds.

Corollary 5.2: Consider a two-dimensional birational mapping with rational first integral $R = P/Q$. Consider also the irreducible components of the singular set of the first type $S_I^{(j)}(\mathbf{f})$ for which the singularities are globally confined. Then the maximum of the degrees of P and Q is bounded below by the number

$$\sum_j (k'_j - 1), \tag{5.7}$$

where k'_j is given by the lowest value k_j of Theorem 3.1.

In example (2.2), the singular set has a unique component in which singularities are globally confined with $k = 4$ and the degree of the first integral is exactly $k - 1 = 3$. Note, however, that the corollary only provides a lower bound as shown in the following example,

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} -y_n + x_n + \frac{a}{x_n} \\ x_n \end{pmatrix}, \tag{5.8}$$

where a is a nonvanishing complex number. Here again, one can check that singularities of the form $(0, y)$ are confined in four steps and since confinement is global, the lower bound for the numerator and denominator of a first integral is 3. However, the rational first integral $I = P/Q$ with lowest degree possible on P and Q is given by

$$I = (x_n - y_n)^2 (x_n y_n - a)^2, \tag{5.9}$$

which is of degree 6, not 3 (note, however, the rather degenerate nature of the first integral).

Next, we turn our attention to singularities of type II in the case when $S_{II}^{(j)}(\mathbf{f}) \not\subset \bar{L}_{i,c}$ for all c and all i . From the proof of Theorems 3.2 and 4.2, one sees that, as long as the singularities are not confined, the iterates of $S_{II}^{(j)}(\mathbf{f}) \not\subset \bar{L}_c$ must be roots of both P_i and Q_i for all rational first integrals P_i/Q_i . Therefore, we have the following corollary.

Corollary 5.3: Consider an algebraically integrable p -dimensional birational mapping with the rational first integral $R = P/Q$. Consider also an irreducible component of the singular set of the second type $S_{II}^{(j)}(\mathbf{f})$ whose singularities are globally confined. Let k_j be the corresponding confinement number. Then the finite elements of the set

$$\bigcup_{i=1}^{k_j-1} \mathbf{f}^i(S_{II}^{(j)}(\mathbf{f}))$$

are roots of both P and Q .

This corollary is best illustrated in example (2.15) where $(0, a)$ and $(1/a, 0)$ are roots of both the numerator and denominator of the rational function (2.17).

The previous corollary has an immediate consequence for the existence of polynomial first integrals.

Corollary 5.4: Consider an algebraically integrable p -dimensional birational mapping. If this system admits singularities of the second type which are globally confined, then there is no polynomial first integral.

The last result concerns the case when the irreducible component of the singular set lies inside the closure of a level set of the first integral. In Secs. III and IV, it was found that if \mathbf{f}^k (for $k \in \mathbb{N}$) is well-defined on $S^{(j)}(\mathbf{f})$ and its Jacobian is nonzero, then $\mathbf{f}^k(S^{(j)}(\mathbf{f}))$ must be a set of the form of the RHS of (3.8), hence it lies on a level set. This is stated in the following corollary.

Corollary 5.5: Consider a p -dimensional birational mapping $\mathbf{x}_{n+1} = \mathbf{f}(\mathbf{x}_n)$ together with an irreducible component of the singular set $S^{(j)}(\mathbf{f})$ for which the singularities are ubiquitous. If \mathbf{f}^k is well-defined almost everywhere on $S^{(j)}(\mathbf{f})$ and its Jacobian is nonzero, then $\mathbf{f}^k(S^{(j)}(\mathbf{f}))$ lies inside the closure of a level set of the first integrals.

Note that the case $k=0$ should also be included meaning that $S^{(j)}(\mathbf{f})$ itself lies inside the closure of a level set of the first integrals.

This corollary can be applied to example (2.18). Since the two sets of singularities $((x, 0)$ and $(0, y))$ are ubiquitous, they are level sets of the first integral (2.19).

As a final example, consider the system

$$\begin{pmatrix} x_{n+1} \\ y_{n+1} \end{pmatrix} = \mathbf{f}(x_n, y_n) = \begin{pmatrix} -y_n + x_n + \frac{a}{x_n^2} \\ x_n \end{pmatrix}, \tag{5.10}$$

where a is a nonzero complex number. This example was first used in Ref. 17 to show that singularity confinement is not sufficient for integrability. Following example (2.2), it is easy to

show that singularities of the form $(0,y)$ are generically globally confined. However, numerical analysis performed on this system strongly suggests that it exhibits chaotic behavior.¹⁷

VI. CONCLUSIONS

In this article, we defined different types of singularities and their confinement property for autonomous discrete birational dynamical systems. The theory of singularity confinement was developed more than 10 years ago^{11,12,27} and different types of singularities have been implicitly used. Here, in order to relate algebraic integrability to singularity confinement we have found necessary to give a formal definition of different singularities and their confinement property.

In Secs. III and IV, we showed that algebraic integrability for birational mappings implies that singularities are either locally confined or some preimages are not well-defined.

In Sec. V, we studied global properties of birational mappings and showed that there are three types of behavior for singularities compatible with algebraic integrability. Namely, singularities are either globally confined, periodic, or ubiquitous. Remarkably, in the seminal papers describing singularity confinement, similar behaviors were found heuristically to be compatible with integrability.^{11–13,15,27,28} However, the classical approach relies mostly on a local analysis of the singularities. An important aspect of the results shown in this article is that global properties have to be satisfied for a system to be algebraically integrable. To unify these different concepts, we define the confinement property for a discrete system in the following way:

Definition: A birational mapping is said to have the **confinement property** if its singularities are, generically, globally confined, periodic, or ubiquitous.

Then the most important result of this article can be rephrased as follows.

Theorem: An algebraically integrable birational mapping has the confinement property.

In particular, this result shows that every mapping in the QRT-family^{15,26} has the confinement property both in the symmetric and asymmetric cases. This fact also holds for the recently discovered class of integrable birational discrete systems discovered in Ref. 22. Moreover, from examples such as system (5.10), we have strong numerical evidence that the confinement property is necessary but not sufficient for algebraic integrability.

In this article, we also found sufficient conditions for nonintegrability. Essentially, if singularities are confined in one direction and not in the other, the system cannot be algebraically integrable.

Finally, the remaining corollaries of Sec. V provide information on the existence and degree of rational first integrals directly from the singularity confinement procedure.

All the examples of integrable discrete systems considered in this article have a first integral of genus 0 or 1. It is not possible to find a birational mapping of infinite order preserving an algebraic curve of genus 2 or higher. The argument proving this fact is due to Veselov (Ref. 31, p. 35). It makes use of the Hurwitz theorem which tells us that the automorphic group of an algebraic curve of genus 2 or higher is finite.

Another important question concerns the relation between the confinement property and other types of integrability for discrete dynamical systems (see, for example, Refs. 2, 4, 5, 17, and 30). In particular, in a work by Bellon,³ the proof that two-dimensional birational mappings admitting a first integral satisfy the *algebraic entropy*^{4,17} integrability condition is given.

Since the only mappings studied here were birational, we could have made use of the projective space in which the point at infinity plays no particular role. Hence only singularities of type II should be studied. However, the ideas presented in this article could not be extended for the case of nonautonomous or discrete dynamical systems that are not birational. For instance, the integrable nonrational mapping derived in Refs. 18–20 could not be studied. It is therefore crucial to give a definition of singularity confinement that can be applied to general discrete systems.

ACKNOWLEDGMENTS

A.G. acknowledges a Fellowship from the Sloan Foundation. S.L. acknowledges a post-doctoral fellowship from NSERC (National Science and Engineering Research Council of Canada). The authors are grateful to Basil Grammaticos, Alfred Ramani, and John Roberts for helpful discussions.

- ¹Ablowitz, M. J., Ramani, A. and Segur, H., "Nonlinear evolution equations and ordinary differential equations of Painlevé type," *Lett. Nuovo Cimento Soc. Ital. Fis.* **23**, 333–338 (1978).
- ²Ablowitz, M. J., Halburd, R., and Herbst, B., "On the extension of the Painlevé property to difference equations," *Nonlinearity* **13**, 889–905 (2000).
- ³Bellon, M. P., "Algebraic entropy of birational maps with invariant curves," *Lett. Math. Phys.* **50**, 79–90 (1999).
- ⁴Bellon, M. P. and Viallet, C-M., "Algebraic entropy," *Commun. Math. Phys.* **204**, 425–437 (1999).
- ⁵Conte, R. and Musette, M., "A new method to test discrete Painlevé equations," *Phys. Lett. A* **223**, 439–448 (1996).
- ⁶Evans, N. W., "Superintegrability in classical mechanics," *Phys. Rev. A* **41**, 5666–5676 (1990).
- ⁷Fris, J., Mandrosov, V., Smorodinsky, Y. A., Uhler, M., and Winternitz, P., "On higher symmetries in quantum mechanics," *Phys. Lett.* **16**, 354–356 (1965).
- ⁸Gambier, B., "Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes," *Acta Math.* **33**, 1–55 (1910).
- ⁹Goriely, A., *Integrability and Nonintegrability of Dynamical Systems* (World Scientific, Singapore, 2001).
- ¹⁰Goriely, A., "Integrability, partial integrability, and nonintegrability for systems of ordinary differential equations," *J. Math. Phys.* **37**, 1871–1893 (1996).
- ¹¹Grammaticos, B., Ramani, A., and Papageorgiou, V. G., "Do integrable mappings have the Painlevé property?" *Phys. Rev. Lett.* **67**, 1825–1826 (1991).
- ¹²Grammaticos, B., Ramani, A., and Tamizhmani, K. M., "Nonproliferation of pre-images in integrable mappings," *J. Phys. A* **27**, 559–566 (1994).
- ¹³Grammaticos, B. and Ramani, A., "The Gambier mapping," *Physica A* **223**, 125–136 (1995).
- ¹⁴Grammaticos, B., Ramani, A., and Lafortune, S., "The Gambier mapping, revisited," *Physica A* **253**, 260–270 (1998).
- ¹⁵Grammaticos, B., Nijhoff, F., and Ramani, A., "Discrete Painlevé equations," in *The Painlevé Property*, Proceedings, Cargèse 1996, edited by R. Conte, CRM Ser. Math. Phys. (Springer, New York, 1999), pp. 413–516.
- ¹⁶Gravel, S. and Winternitz, P., "Superintegrability with third-order integrals in quantum and classical mechanics," *J. Math. Phys.* **43**, 5902–5912 (2002).
- ¹⁷Hietarinta, J. and Viallet, C-M., "Singularity confinement and chaos in discrete systems," *Phys. Rev. Lett.* **81**, 325–328 (1998).
- ¹⁸Iatrou, A. and Roberts, J. A. G., "Integrable mappings of the plane preserving biquadratic invariant curves," *J. Phys. A* **34**, 6617–6636 (2001).
- ¹⁹Iatrou, A. and Roberts, J. A. G., "Integrable mappings of the plane preserving biquadratic invariant curves II," *Nonlinearity* **15**, 459–489 (2002).
- ²⁰Iatrou, A. and Roberts, J. A. G., "Integrable mappings of the plane preserving biquadratic invariant curves III," *Physica A* **326**, 400–411 (2003).
- ²¹Ince, E. L., *Ordinary Differential Equations* (Dover, New York, 1956).
- ²²Kimura, K., Yahagi, H., Hirota, R., Ramani, A., Grammaticos, and Y. Ohta, "A new class of integrable discrete systems," *J. Phys. A* **35**, 9205–9212 (2002).
- ²³Kummer, M., Churchill, R. C., and Rod, D. L., "On a result of Brun," *Can. Math. Bull.* **33**, 175–180 (1990).
- ²⁴Olver, P. J., *Applications of Lie Groups to Differential Equations* (Springer-Verlag, New York, 1993).
- ²⁵Painlevé, P., *Oeuvres de Paul Painlevé. Tome I* (Editions du CNRS, Paris, 1973).
- ²⁶Quispel, G. R. W., Roberts, J. A. G., and Thompson, C. J., "Integrable mappings and soliton equations II," *Physica D* **34**, 183–192 (1989).
- ²⁷Ramani, A., Grammaticos, B., and Hietarinta, J., "Discrete versions of the Painlevé equations," *Phys. Rev. Lett.* **67**, 1829–1832 (1991).
- ²⁸Ramani, A., Grammaticos, B., and Karra, G., "Linearizable mappings," *Physica A* **180**, 115–127 (1992).
- ²⁹Ramani, A., Grammaticos, B., Tamizhmani, K. M., and Lafortune, S., "Again, linearizable mappings," *Physica A* **252**, 138–150 (1998).
- ³⁰Roberts, J. A. G., Jogia, D., and Vivaldi, F., "The Hasse-Weil Bound and Integrability Detection in Rational Maps," *J. Nonlinear Math. Phys.* **10**, Suppl. 2, 166–180 (2003).
- ³¹Veselov, A. P., "Integrable maps," *Russ. Math. Surveys* **46**, 1–51 (1991).
- ³²Yoshida, H., "Necessary conditions for the existence of algebraic first integrals I," *Celest. Mech.* **31**, 363–379 (1983).
- ³³Yoshida, H., "Necessary conditions for the existence of algebraic first integrals II," *Celest. Mech.* **31**, 381–399 (1983).